

the treatment of cognitive impairments in people with schizophrenia[J]. Am J Psychiatry,2008 ,165(1) :82289.

- [4] Lee BJ ,Lee J G , Kim YH. A 122week ,double2blind ,placebo2 controlled trial of donepezil as an adjunct to haloperidol for treating cognitive impairments in patients with chronic schizophrenia[J].J Psychopharmacol ,2007 ,21(4) :4212427.
- [5] Akhondzadeh S , Gerami M , Noroozian M , et al. A 122week , double2blind , placebo2controlled trial of donepezil adjunctive treatment to risperidone in chronic and stable schizophrenia [J]. Prog Neuropsychopharmacol Biol Psychiatry , 2008 , 32 (8) :181021815.
- [6] Keefe RS ,Malhotra A K ,Meltzer HY , et al. Efficacy and safety of donepezil in patients with schizophrenia or schizoaffective disorder : significant placebo/ practice effects in a 122week , randomized , double2blind , placebo2controlled trial [J]. Neuropsychopharmacology ,2008 ,33(6) :121721228.
- [7] Rosse

域、中央的 Rho 结合的卷曲螺旋结构域和羧基端 p H 结构域。
在静息状

www.cnki.net

还可通过介导缺氧或凝血酶引起的内皮型 NOS (endo helial NOS,eNOS) 表达下调^[21],使 NO 扩张血管增加脑缺血区血流量、抑制血小板聚集、抑制小血管缺血后炎症反应的脑保护作用下降。有研究发现大鼠大脑中动脉阻塞后,出现三磷酸腺苷(ATP)耗竭,缺血造成细胞内钙离子增多,引发级联反应,与迟发性细胞死亡或凋亡有关,应用 Fausdil 在各时段增加局部脑血流量,可延缓缺血脑组织进入不可逆的细胞死亡过程,体现脑保护作用^[22]。Rikitake 等^[23]在大脑中动脉闭塞大鼠中应用不同剂量 Fasudil 抑制 Rho 激酶活性,结果显示,内皮型一氧化氮酶 mRNA 及其蛋白表达随 Fausdil 浓度的增加而增加,相应的 eNOS 活性和 NO 含量也增加,从而使脑梗死体积较未治疗组缩小 33%,神经功能评分提高 37%。而在 eNOS 缺陷小鼠中未能观察到神经保护作用。



www.cnki.net

288.6 566.5 456 527.2 m 1 195.7 522.3 1 194.3 521.9161.5 689.4 Td (0) Tj ET